

PRODUCTS TO AVOID PRIOR TO YOUR PROCEDURE
(These are drugs that can thin your blood and may cause increased bleeding)
Avoid these medications for at least 5 days before your procedure.

Advil	Motrin
Aleve	Naprosyn
Alka Seltzer	Naprelan
Anaprox	Naproxen
Anacin	Norgesic
Arthritis Pain Formula	Norwich
Ascriptin	Nuprin
Aspergum	P-A-C Analgesic
Aspirin	Pepto- Bismal
Azoid	Percodan
BC Powders	Plavix
Bextra	Robaxisal
Bufferin	Salflex
Carna Arthritis Medication	Salsalate
Celebrex	Salsitab
Clinoril	Sine-Aid
Coumadin	Soma Compound
Darvon Compound / Darvocet	Stanback
Disalcid	Synalgos-DC
Doan's Pills	Talwin
Dristan	Therapy Bayer Caplets
Easpirin	Tolectin
Ecotrin	Tolmetin
Effervescent Tablets	Toradol
Empirin	Trigesic
Equagesic	Trilisate
Etodolac	Urisinus-Inlay
Excedrin	Vanquish
Feldene	Vioxx
Fenoprex	Warfarin
Fiorinal	
Formula Caplets	
Goody Powders	
Haltran	
Indocin	
Ibuprofen	
Ibu-Tab	
Lodine	
Loritab ASA	
Magsal	
Midol	
Mobic	
Mobigesic	
Monogesic Tablets	